

## INDEX TO VOLUME 73

## AUTHORS

- Allen, Sarah G.: see Webber and Allen, 60-61.
- Asay, Christopher E.: Habitat and Productivity of Cooper's Hawks Nesting in California, 80-87.
- Ault, Jerald S., and John D. DeMartini: Movement and Dispersion of Red Abalone, *Haliotis rufescens*, in Northern California, 196-213.
- Brooks, Andrew J.: Two Species of Kyphosidae Seen in King Harbor, Redondo Beach, California, 49-50.
- Coblentz, Bruce E.: Yellowtail Chafing on a Shark: Parasite Removal?, 244.
- Cole, P.J.: see Crawford, Cole, Kilbride, and Fairbrother, 245-247.
- Compagno, Leonard J.V.: see Ebert, Compagno, and Natanson, 117-123.
- Crawford, J.A., P.J. Cole, K.M. Kilbride, and A. Fairbrother: Atypical Plumage of a Female California Quail, 245-247.
- DeMartini, John D.: see Ault and DeMartini, 196-213.
- Dole, Jim W.: see Perry, Dole, and Holl, 156-162.
- Duhamel, G.E.: see Kent, Duhamel, Foott, and Hedrick, 99-105. Ebert, David A., Leonard J.V. Compagno, and Lisa J. Natanson: Biological Notes on the Pacific Sleeper Shark, *Somniosus pacificus* (Chondrichthyes: Squalidae), 117-123.
- Erickson, Daniel L. and Ellen K. Pikitch: First Oregon Record for the Cowcod, *Sebastes levis*, 192.
- Euliss, Ned H. Jr., and Gail Grodhaus: Management of Midges and Other Invertebrates for Waterfowl Wintering in California, 238-243.
- Fairbrother, A.: see Crawford, Cole, Kilbride, and Fairbrother, 245-247.
- Foott, J.S.: see Kent, Duhamel, Foott, and Hedrick, 99-105.
- Fritzsche, Ronald A.: see Ward and Fritzsche, 175-187.
- Gilbert, Barrie K.: see Hastings and Gilbert, 188-191.
- Gotshall, Daniel W.: The use of Baited Stations by Divers to Obtain Fish Relative Abundance Data, 214-229.
- Grodhaus, Gail: see Euliss and Grodhaus, 238-243.
- Hartmann, A. Rucker: Movement of Scorpionfishes (Scorpaenidae: *Sebastes* and *Scorpaena*) in the Southern California Bight, 68-79.
- Hastings, Bruce C., and Barrie K. Gilbert: Extent of Human-Bear Interactions in the Backcountry of Yosemite National Park, 188-191.
- Hedrick, R.P.: see Kent, Duhamel, Foott, and Hedrick, 99-105.
- Hemmer, M.J.: see Russell, Middaugh, and Hemmer, 169-174.
- Holl, Stephan A.: see Perry, Dole, and Holl, 156-162.
- Ivey, Gary L.: Winter Foods of American Coots in the Northern San Joaquin Valley, California, 45-48.
- Kent, M.L., G.E. Duhamel, J.S. Foott, and R.P. Hedrick: Chronic Branchitis (Hamburger Gill Disease) of Channel Catfish in California and its Possible Mysosporean Etiology, 99-105.
- Kilbride, K.M.: see Crawford, Cole, Kilbride, and Fairbrother, 245-247.
- Knutson, Arthur C. Jr.: Comparative Catches of Ocean Sport-Caught Salmon using Barbed and Barbless Hooks and Estimated 1984 San Francisco Bay Area Charterboat Shaker Catch, 106-116.
- Lea, Robert N., and Richard H. Rosenblatt: Occurrence of the Family Notacanthidae (Pisces) from Marine Waters of California, 51-53.
- Lea, Robert N.: On the Second Record of *Barbourisia rufa*, the Velvet Whalefish, from California, 124.
- Marshall, William H.: see Miller and Marshall, 37-44.
- Middaugh, D.P.: see Russell, Middaugh, and Hemmer, 169-174.
- Miller, Kathy Ann, and William H. Marshall: Food Habits of Large Monkeyface Prickleback, *Cebidichthys violaceus*, 37-44.
- Natanson, Lisa J.: see Ebert, Compagno, and Natanson, 117-123.
- Perry, William M., Jim W. Dole, and Stephen A. Holl: Analysis of the Diets of Mountain Sheep from the San Gabriel Mountains, California, 156-162.
- Pikitch, Ellen K.: see Erickson and Pikitch, 192.
- Randall, John E.: Refutation of Lengths of 11.3, 9.0, and 6.4 m Attributed to the White Shark, *Carcharodon carcharias*, 1635-168.
- Reilly, Paul N.: Population Studies of Rock Crabs, *Cancer antennarius*, Yellow Crabs, *C. anthonyi*, and Kellet's Whelks, *Kelletia kelletii*, in the Vicinity of a Proposed Liquefied Natural Gas Terminal at Little Cojo Bay, Santa Barbara County, California, 88-98.
- Rienecker, Warren C.: Population Trends, Distribution, and Survival of Canada Geese in California and Western Nevada, 1949-79, 21-36.

## INDEX TO VOLUME 73

## AUTHORS

- Allen, Sarah G.: see Webber and Allen, 60-61.
- Asay, Christopher E.: Habitat and Productivity of Cooper's Hawks Nesting in California, 80-87.
- Ault, Jerald S., and John D. DeMartini: Movement and Dispersion of Red Abalone, *Haliotis rufescens*, in Northern California, 196-213.
- Brooks, Andrew J.: Two Species of Kyphosidae Seen in King Harbor, Redondo Beach, California, 49-50.
- Coblentz, Bruce E.: Yellowtail Chafing on a Shark: Parasite Removal?, 244.
- Cole, P.J.: see Crawford, Cole, Kilbride, and Fairbrother, 245-247.
- Compagno, Leonard J.V.: see Ebert, Compagno, and Natanson, 117-123.
- Crawford, J.A., P.J. Cole, K.M. Kilbride, and A. Fairbrother: Atypical Plumage of a Female California Quail, 245-247.
- DeMartini, John D.: see Ault and DeMartini, 196-213.
- Dole, Jim W.: see Perry, Dole, and Holl, 156-162.
- Duhamel, G.E.: see Kent, Duhamel, Foott, and Hedrick, 99-105. Ebert, David A., Leonard J.V. Compagno, and Lisa J. Natanson: Biological Notes on the Pacific Sleeper Shark, *Somniosus pacificus* (Chondrichthyes: Squalidae), 117-123.
- Erickson, Daniel L. and Ellen K. Pikitch: First Oregon Record for the Cowcod, *Sebastes levis*, 192.
- Euliss, Ned H. Jr., and Gail Grodhaus: Management of Midges and Other Invertebrates for Waterfowl Wintering in California, 238-243.
- Fairbrother, A.: see Crawford, Cole, Kilbride, and Fairbrother, 245-247.
- Foott, J.S.: see Kent, Duhamel, Foott, and Hedrick, 99-105.
- Fritzsche, Ronald A.: see Ward and Fritzsche, 175-187.
- Gilbert, Barrie K.: see Hastings and Gilbert, 188-191.
- Gotshall, Daniel W.: The use of Baited Stations by Divers to Obtain Fish Relative Abundance Data, 214-229.
- Grodhaus, Gail: see Euliss and Grodhaus, 238-243.
- Hartmann, A. Rucker: Movement of Scorpionfishes (Scorpaenidae: *Sebastes* and *Scorpaena*) in the Southern California Bight, 68-79.
- Hastings, Bruce C., and Barrie K. Gilbert: Extent of Human-Bear Interactions in the Backcountry of Yosemite National Park, 188-191.
- Hedrick, R.P.: see Kent, Duhamel, Foott, and Hedrick, 99-105.
- Hemmer, M.J.: see Russell, Middaugh, and Hemmer, 169-174.
- Holl, Stephan A.: see Perry, Dole, and Holl, 156-162.
- Ivey, Gary L.: Winter Foods of American Coots in the Northern San Joaquin Valley, California, 45-48.
- Kent, M.L., G.E. Duhamel, J.S. Foott, and R.P. Hedrick: Chronic Branchitis (Hamburger Gill Disease) of Channel Catfish in California and its Possible Mysosporean Etiology, 99-105.
- Kilbride, K.M.: see Crawford, Cole, Kilbride, and Fairbrother, 245-247.
- Knutson, Arthur C. Jr.: Comparative Catches of Ocean Sport-Caught Salmon using Barbed and Barbless Hooks and Estimated 1984 San Francisco Bay Area Charterboat Shaker Catch, 106-116.
- Lea, Robert N., and Richard H. Rosenblatt: Occurrence of the Family Notacanthidae (Pisces) from Marine Waters of California, 51-53.
- Lea, Robert N.: On the Second Record of *Barbourisia rufa*, the Velvet Whalefish, from California, 124.
- Marshall, William H.: see Miller and Marshall, 37-44.
- Middaugh, D.P.: see Russell, Middaugh, and Hemmer, 169-174.
- Miller, Kathy Ann, and William H. Marshall: Food Habits of Large Monkeyface Prickleback, *Cebidichthys violaceus*, 37-44.
- Natanson, Lisa J.: see Ebert, Compagno, and Natanson, 117-123.
- Perry, William M., Jim W. Dole, and Stephen A. Holl: Analysis of the Diets of Mountain Sheep from the San Gabriel Mountains, California, 156-162.
- Pikitch, Ellen K.: see Erickson and Pikitch, 192.
- Randall, John E.: Refutation of Lengths of 11.3, 9.0, and 6.4 m Attributed to the White Shark, *Carcharodon carcharias*, 1635-168.
- Reilly, Paul N.: Population Studies of Rock Crabs, *Cancer antennarius*, Yellow Crabs, *C. anthonyi*, and Kellet's Whelks, *Kelletia kelletii*, in the Vicinity of a Proposed Liquefied Natural Gas Terminal at Little Cojo Bay, Santa Barbara County, California, 88-98.
- Rienecker, Warren C.: Population Trends, Distribution, and Survival of Canada Geese in California and Western Nevada, 1949-79, 21-36.

- Rienecker, Warren C.: Migration and Distribution of Northern Pintails Banded in California, 139-155.
- Rienecker, Warren C.: Survival and Recovery Rate Estimates of Northern Pintails Banded in California, 1946-79, 230-237.
- Rosenblatt, Richard H.: see Lea and Rosenblatt, 51-53.
- Russell, G. A., D. P. Middaugh, M. J. Hemmer.: Reproductive Rhythmicity of the Atherinid Fish, *Colpichthys regis*, from Estero del Soldado, Sonora, Mexico, 169-174.
- Seigel, Jeffrey A.: Record of the Twinpored Eel, *Xenomystax atrarius* (Anguilliformes: Congridae) from California Waters, 57-59.
- Spratt, Jerome D.: Variation in the Growth Rate of Pacific Herring from San Francisco Bay, California, 132-138.
- Ward, David L., and Ronald A. Fritzsche: Comparison of Meristic and Morphometric Characters among and Within Subspecies of the Sacramento Sucker (*Catostomus occidentalis*) Ayres, 175-187.
- Warner, Ronald W.: Age and Growth of Male Dungeness Crabs, *Cancer magister*, in Northern California, 4-20.
- Webber, Marc A., and Sarah G. Allen: Resightings of Two Rehabilitated and Released Harbor Seals in California, 60-61.
- Wicksten, Mary K.: Range Extensions of Offshore Decapod Crustaceans from California and Western Mexico, 54-56.

## SUBJECT

- Abalone, red: Movement and dispersion of, in northern California, 196-213
- Baited stations: Use of, by divers to obtain fish relative abundance data, 214-229
- Bear-human interactions: In the backcountry of Yosemite National Park, 188-191
- Branchitis, chronic: Of channel catfish, 99-105
- Coot, American: Winter foods of, 45-48
- Cowcod: First Oregon record, 192
- Crab, Dungeness: Age and growth of males, 4-20
- Crab, rock: Population studies of, 88-98
- Crab, yellow: Population studies of, 88-98
- Crustaceans, decapod: Range extensions of, 54-56
- Eel, twinpored: Record from California waters, 57-59
- El Niño: 50, 113, 134
- Food habits: Of large monkeyface prickleback, 37-44
- Foods, winter: Of American coots, 45-48
- Geese, Canada: Population trends, distribution, and survival, 21-36
- Gill disease, hamburger: Of channel catfish, 99-105
- Hawk, Cooper's: Habitat and productivity of, 80-87
- Herring, Pacific: Variation in the growth rate, 132-138
- Kyphosidae: Two species seen in King Harbor, Redondo Beach, 49-50
- Midges: Management of, for waterfowl wintering in California, 238-243
- Notacanthidae: Occurrence from marine waters of California, 51-53
- Pintail, northern: Migration and distribution of, banded in California, 139-155
- Pintail, northern: Survival and recovery rate estimates of, banded in California, 230-237
- Population studies: Of rock crabs, yellow crabs, and Kellet's whelks, 88-98
- Population trends: Of Canada geese in California and western Nevada, 21-36
- Prickleback, monkeyface: Food habits of, 37-44
- Quail, California: Atypical plumage, 245-247
- Range extension: Of offshore decapod crustaceans, 54-56
- Reproductive rhythmicity: Of the atherinid fish, *Colpichthys regis*, 169-174
- Salmon, ocean sport-caught: Comparative catches of, using barbed and barbless hooks, 106-116
- Scorpion fishes: Movement in the Southern California Bight, 68-79
- Seal, harbor: Resightings of two rehabilitated and released, 60-61

- Shark, Pacific sleeper: Biological notes on, 117-123  
 Shark: Yellowtail chafing on, 244  
 Shark, white: Refutation of lengths of 11.3, 9.0, and 6.4 m, 163-168  
 Sheep, mountain: Analysis of the diets of, 156-162  
 Sucker, Sacramento: Comparison of meristic and morphometric characters among and within subspecies, 175-187  
 Waterfowl: Management of midges and other invertebrates for, 238-243  
 Whalefish, velvet: On the second record of, from California, 124  
 Whelk, Kellet's: Population studies of, 88-98  
 Yellowtail: Chafing on a shark, 244

## SCIENTIFIC NAMES

- Accipiter cooperii*: 80-87  
*Adenostoma fasciculatum*: 160  
*Aglaophenia* sp.: 41  
*Ahnfeltia plicata*: 39  
*Anas acuta acuta*: 238  
*Anas acuta*: 46, 139-155  
*Anas americana*: 26, 233  
*Anas crecca carolinensis*: 238  
*Anas platyrhynchos*: 35, 230-237, 238  
*Anisotremis davidsonii*: 49  
*Anoplopoma fimbria*: 124  
*Anser americana*: 147  
*Anser caerulescens caerulescens*: 26, 147  
*Anser crecca carolinensis*: 147  
*Apedium subcinctum*: 239  
*Aplidium* sp.: 41  
*Archidistoma* sp.: 41  
*Archidistoma ritteri*: 41  
*Arctostaphylos* sp.: 160  
*Aster alexis*: 47  
*Atherinops*: 169, 170  
*Athya collaris*: 233  
*Avicennia germinans*: 170  
Axiidae: 54  
*Barbourisia rufa*: 124  
Barbourisiidae: 124  
Bothidae: 220  
*Botryoglossum farlowianum*: 38, 41  
*Branta canadensis maxima*: 34  
*Branta canadensis minima*: 35  
*Branta canadensis moffitti*: 21-36  
*Branta canadensis occidentalis*: 29  
*Callianassa goniophthalma*: 55  
Callianassidae: 55  
*Calliarthron tuberculosum*: 200  
*Callipepla californica*: 245  
*Callophyllis crenulata*: 39  
*Callophyllis pinnata*: 38  
*Callophyllis violacea*: 38  
*Calocaris quinqueseriatus*: 54  
*Cancer antennarius*: 88-98  
*Cancer anthonyi*: 88-98  
*Cancer magister*: 4-20  
*Cancer productus*: 96  
Carangidae: 244  
*Carassius auratus*: 241  
*Carcharodon carcharias*: 163-168  
*Castilleja affinis*: 159  
*Catostomus occidentalis*: 175-187  
*Catostomus occidentalis humboldtianus*: 175-187  
*Catostomus occidentalis lacusanserinus*: 175-187  
*Catostomus occidentalis mniotiltus*: 175-187  
*Catostomus occidentalis occidentalis*: 175-187  
*Ceanothus crassifolius*: 160  
*Ceanothus leucodermis*: 159  
*Cebidichthys violaceus*: 37-44  
*Centrocercus clavulatus*: 39  
*Cephaloscyllium ventriosum*: 220  
*Ceramium* sp.: 39  
*Cercocarpus betuloides*: 158, 159  
*Cercocarpus ledifolius*: 159, 161  
*Chione*: 170  
Chironomidae: 46, 47, 239  
*Chironomus decorus*: 239  
*Chironomus stigmaterus*: 239  
Ciccadellidae: 47  
*Cladophora columbiana*: 38  
*Clupea harengus*: 69  
*Clupea harengus pallasii*: 132-138  
*Colinus virginianus*: 245  
*Colpichthys regis*: 169-174  
Congridae: 57, 59  
*Corallina vancouveriensis*: 39  
Corixidae: 47  
*Costia*: 102  
Cottidae: 220  
*Coturnix coturnix japonica*: 245  
*Cricotopus* sp.: 239  
*Cryptopleura corallinara*: 39  
*Cryptopleura lobulifera*: 38  
*Cryptopleura violacea*: 38, 41  
*Cryptosiphonia woodii*: 38  
*Cyprinus carpio*: 241  
Decapoda: 54-56

- Dilsea californica*: 39  
*Distaplia occidentalis*: 41  
*Distaplia* sp.: 41  
*Distichlis spicata*: 47  
*Dodecaceria concharum*: 199  
Dytiscidae: 47  
*Echinochloa crusgalli*: 47, 239  
*Echinodorus cordifolius*: 239  
*Embiotoca jacksoni*: 217  
Embiotocidae: 220  
*Encelia californica*: 159  
*Endocladia muricata*: 39  
*Engraulis mordax*: 107, 136  
*Enteromorpha intestinalis*: 39  
*Enteromorpha linza*: 39  
Ephydriidae: 47  
*Eriodictyon crassifolium*: 160  
*Eriogonum fasciculatum*: 158, 159  
*Eriogonum ovalifolium*: 161  
*Eriogonum umbellatum*: 161  
*Eriophyllum confertiflorum*: 159  
*Erythrophyllum delesserioides*: 39  
*Farlowia mollis*: 39  
*Fulca americana*: 45-48  
*Galathea californiensis*: 55  
Galatheididae: 55  
*Gambusia affinis*: 241  
*Garrya veatchii*: 158, 159  
*Gasterosteus aculeatus*: 241  
*Gastroclonium coulteri*: 38, 41  
*Celidium coulteri*: 39  
*Celidium purpurascens*: 39  
*Gigartina canaliculata*: 39  
*Gigartina corymbifera*: 39  
*Gigartina volans*: 39  
*Gracilaria sjoestedtii*: 39  
*Gymnogongrus linearis*: 39  
*Haliotis corrugata*: 211, 212  
*Haliotis iris*: 210  
*Haliotis midae*: 210  
*Haliotis ruber*: 211  
*Haliotis rufescens*: 196-213  
*Haliotis sorenseni*: 211  
*Haliotis tuberculata*: 211  
*Halymenia californica*: 39  
*Heleochoa shoenoides*: 47  
*Henneguya*: 99-105  
*Henneguya exilis*: 100, 103  
*Heptacarpus* sp.: 41  
*Hermosilla azurea*: 49  
*Heteromeles arbutifolia*: 159  
Hexagrammidae: 220  
Hippolytidae: 54  
*Hymenena flabelligera*: 38, 41  
*Hypsurus caryi*: 217  
*Ictalurus nebulosus*: 100  
*Ictalurus punctatus*: 99-105  
*Idotea stenops*: 41  
*Iridaea cordata*: 38, 39, 41, 42  
*Iridaea flaccida*: 39  
Istiophoridae: 244  
*Juncus* sp.: 47  
*Kelletia kelletii*: 88-98  
Kyphosidae: 49-50  
*Kyphosus analogus*: 49-50  
Labridae: 220  
*Leptodactylon californicum*: 159  
*Leuresthes sardina*: 169, 174  
*Leuresthes tenuis*: 169, 174  
Lithodidae: 55  
*Loligo opalescens*: 4  
Lymnaeidae: 47  
*Macdonaldia challengerii*: 51  
*Macrocystis pyrifera*: 89, 211, 215  
*Mastocarpus jordinii*: 39  
*Mastocarpus papillatus*: 38, 41  
*Menidia menidia*: 169, 172  
*Mentzelia laevicaulis*: 159  
*Merluccius productus*: 57  
*Microcladia borealis*: 38  
*Microcladia coulteri*: 39  
*Microstomus pacificus*: 51, 124  
Moblidae: 244  
*Morone saxatilis*: 115  
*Munida hispida*: 55  
Muraenesocidae: 59  
*Najas guadalupensis*: 239  
*Neoclinus* spp.: 220  
*Neoclinus uninotatus*: 217  
*Neoptilota californica*: 39  
*Neoptilota densa*: 39  
*Neoptilota hypnoides*: 39  
Nettastomidae: 59  
*Nienburgia andersonii*: 39  
Notacanthidae: 51-53  
*Notacanthus chemnitzii*: 51-53  
*Octopoteuthis deletron*: 121  
*Odonthalia floccosa*: 39  
*Oncorhynchus kisutch*: 106-116  
*Oncorhynchus tshawytscha*: 106-116, 122  
*Ophiodon elongatus*: 69, 72  
*Ovis canadensis nelsoni*: 156-162  
*Oxyjulis californica*: 227  
Pandalidae: 54  
*Pandalopsis ampla*: 54  
*Paralabrax clathratus*: 217, 220  
*Paralithodes rathbuni*: 55  
Parapaguridae: 55

- Parapagurus haigae*: 55  
*Paspalum distichum*: 47  
 Phasianidae: 245  
*Phasianus colchicus*: 245  
*Phocavulina*: 122  
*Phoca vitulina richardsi*: 60-61  
*Phyllospadix torreyi*: 38  
*Pikea californica*: 39  
*Pinus sabiniana*: 82  
*Pithophora* sp.: 46, 47  
*Platanus racemosa*: 82  
 Pleuronectidae: 220  
*Plocomium cartilagineum*: 38  
*Pogonophorella californica*: 39  
*Polyacanthonotus challengerii*: 51  
*Polycheria* sp.: 41  
*Polyneura latissima*: 38  
*Polysiphonia* sp.: 39  
*Populus fremontii*: 82  
*Porichthys* spp.: 220  
*Porphyra lanceolata*: 39  
*Porphyra perforata*: 38, 40, 41  
*Potamogeton pectinatus*: 239  
*Prionace glauca*: 244  
*Prionitis lanceolata*: 39  
*Prionitis lyallii*: 38  
*Prunus ilicifolia*: 158, 159  
*Psychrolutes phrictus*: 51  
*Pterosiphonia dendroidea*: 39  
*Pterygophora californica*: 95, 215  
*Ptilota filicina*: 38  
*Pycnopia helianthoides*: 199  
*Quercus agrifolia*: 82  
*Quercus douglasii*: 82  
*Quercus lobata*: 82  
*Quercus wislizenii*: 82  
*Rathbunella hypoplecta*: 217, 220  
*Rhamnus californica*: 159  
*Rhamnus crocea*: 159  
*Rhizophora mangle*: 170  
*Rhodoglossum affine*: 38  
*Rhodoglossum roseum*: 38  
*Rhodomela larix*: 39  
*Rhodymenia californica*: 39  
*Rhus ovata*: 159  
*Rumex crispus*: 47  
*Ruppia maritima*: 239  
*Salvia apiana*: 158, 159  
*Sarcodiotheca gaudichaudii*: 39  
*Schizymenia pacifica*: 39  
*Scirpus acutus*: 239  
*Scirpus maritimus*: 47  
*Scirpus* spp.: 47  
*Scorpaena guttata*: 69, 72  
*Scorpaena*: 68-79  
*Scorpaenichthys marmoratus*: 69, 72  
*Scorpaenidae*: 68-79  
*Sebastes*: 68-79  
*Sebastes* spp.: 4  
*Sebastes atrovirens*: 72  
*Sebastes auriculatus*: 72  
*Sebastes carnatus*: 72  
*Sebastes caurinus*: 72  
*Sebastes chlorostictus*: 72  
*Sebastes chrysomelas*: 72  
*Sebastes constellatus*: 72  
*Sebastes dalli*: 72  
*Sebastes entomelas*: 69, 72  
*Sebastes flavidus*: 72  
*Sebastes goodei*: 69  
*Sebastes hopkinsi*: 72  
*Sebastes jordani*: 69  
*Sebastes levis*: 69, 192  
*Sebastes melanops*: 72  
*Sebastes miniatus*: 69, 72  
*Sebastes mystinus*: 72, 137  
*Sebastes nebulosus*: 72  
*Sebastes ovalis*: 72  
*Sebastes paucispinis*: 69, 72  
*Sebastes pinniger*: 72  
*Sebastes rastrelliger*: 72  
*Sebastes rosaceus*: 72  
*Sebastes ruberrimus*: 72  
*Sebastes rubrivinctus*: 72  
*Sebastes serranoides*: 72  
*Sebastes serripes*: 72  
*Sebastes umbrosus*: 72  
*Sebastolobus alascanus*: 121, 124  
*Sectator ocyurus*: 49-50  
*Seriola lalandi*: 244  
*Somniosus microcephalus*: 118  
*Somniosus pacificus*: 117-123  
*Spirontocaris sica*: 54  
 Squalidae: 117  
*Squatina californica*: 220  
*Stenogramme interrupta*: 39  
*Strongylocentrotus*: 215  
*Synodus lucioceps*: 220  
*Thaleichthys pacificus*: 61  
*Thunnus alalunga*: 121

*Trichodina*: 102

*Trichophyra*: 102

*Trifolium* sp.: 47

*Typha latifolia*: 239

*Ulva lobata*: 38, 40, 41, 43

*Ulva taeniata*: 38

*Ursus americanus*: 188-191

*Ursus arctos*: 188

*Viola purpurea*: 159

*Zenomystax atrarius*: 57-59

*Yucca whipplei*: 159

*Zannichellia palustris*: 239